

Furosemide

10mg/ml Solution for injection /
infusion

bowmed
ibisqus

Furosemide is a loop diuretic commonly prescribed to reduce fluid accumulation and manage severe hypertension effectively.

Therapeutic indications

Furosemide 10mg/ml solution for injection/infusion is indicated when no adequate diuresis is achieved with oral administration of furosemide or when oral use is not possible:

- oedemas and/or ascites secondary to cardiac or hepatic disorders
- oedemas secondary to renal disorders
- pulmonary oedema (e.g. in acute heart failure)
- hypertensive crisis (in addition to other therapeutic measures)

Method of administration:

- Intravenous or intramuscular administration.
- As a rule, Furosemide 10 mg/ml solution for injection/infusion is administered intravenously.
- In exceptional cases where neither oral nor intravenous administration is possible, Furosemide 10mg/ml solution for injection/infusion can be administered intramuscularly, but not in acute situations (e.g. not in pulmonary oedema) and not at higher doses.



Strength 20mg/2ml (10mg/ml) Amp x 5 pk

Furosemide

Product Code	BOW0163
EAN Code	5 060130131697
AAH Code	FUR0136B
PL Number	PL 29472/0005
MA Holder	S.A.L.F. S.p.A. Laboratorio Farmacologico
BNF Section	CARDIOVASCULAR SYSTEM
Product Group	Diuretics

Key points



Sold in amber glass ampoules. Keep in the outer carton in order to protect from light.



After intravenous administration of furosemide, onset of action can be expected within 2 to 15 minutes.



Unopened ampoules have a 2 year shelf life.

For more information

bowmed.com
+44 (0)845 6436 703
info@bowmed.com

Bowmed Ibisqus Limited, The Old Dairy,
Brynkinalt Business Centre, Brynkinalt,
Chirk, Wrexham LL14 5NS

Registered Office: Christian Douglass, 2 Jordan Street, Knott Mill, Manchester M15 4PY
Company Reg No. 6204088 Vat No. GB 918 8607 86

Scan or click for:

Summary of Product Characteristics (SPC)
Patient Information Leaflet (PIL)

